

All correspondence to:

The Breastfeeding Network
PO Box 11126, Paisley PA2 8YB
Admin Tel: 0844 412 0995
e-mail: admin@breastfeedingnetwork.org.uk
www.breastfeedingnetwork.org.uk

Colonoscopy and Breastfeeding Mothers

The information provided is taken from various reference sources. It is provided as a guideline. No responsibility can be taken by the author or the Breastfeeding Network for the way in which the information is used. Clinical decisions remain the responsibility of medical and breastfeeding practitioners. The data presented here is intended to provide some immediate information but cannot replace input from professionals.

Colonoscopy is carried out under light sedation in order to examine the inside of the colon. A small flexible tube is passed in through the anus and into the intestine. It may be used to diagnose or exclude conditions such as inflammatory bowel disease or diverticulitis. The examination takes under half an hour and in most cases the patient is not asleep but drowsy and relaxed. For further information see www.patient.co.uk/health/colonoscopy

The colonoscopy procedure is uncomfortable but not painful and is over in a short period. In many instances the preparation period is worse as the urgency to pass motions can be extreme. If a breastfeeding mother needs to have this procedure it may be important that someone is with her to look after the baby during the bowel preparation period, as well as on the day following. She may be drowsy afterwards from lack of food in addition to the colonoscopy and sedation.

In order that the medical staff can examine the gut thoroughly, bowel cleansing is necessary to clear out all faeces in advance of the procedure. This requires a low residue diet for one or two days and the use of a strong laxative the day before the procedure to produce liquid motions as the gut is cleared.

The laxative normally used is a stimulant laxative such as sodium picosulphate (Picolax[®]) which is not known to be absorbed into breastmilk or the osmotic laxative macrogol 3350 (Klean Prep[®]). Macrogol is an inert substance that passes through the gut without being absorbed into the body. It works because it causes water to be retained in the bowel instead of being absorbed into the body. Enemas such as Fleet[®] may also be used without affecting the breastfed baby.

Care should be taken to ensure that the mother remains adequately hydrated throughout the preparation period and on the day of the procedure by drinking plenty of water as permitted by the pre-operative instructions.

Short acting hypnotics such as midazolam, fentanyl and pethidine are used during the procedure itself to sedate the patient and are not contra- indicated during breastfeeding unless the baby is at risk of respiratory depression.

To speak to a Breastfeeding Supporter call the National Breastfeeding Helpline 0300 100 0212

Calls to 0300 numbers cost no more than calls to UK numbers starting 01 and 02 and will be part of any inclusive minutes that apply to your provider and call package

The Breastfeeding Network is a Company Limited by Guarantee Registered in Scotland Company No. 330639
Registered office Whitelaw Wells, 9 Ainslie Place, Edinburgh, EH3 6AT
The Breastfeeding Network is a Registered Scottish Charity No SC027007

©Wendy Jones PhD, MRPharmS and the Breastfeeding Network May 2017

To find your nearest Breastfeeding Supporter call the **Supporterline 0300 100 0210**

Midazolam : The small amounts of **midazolam** excreted into breastmilk would not be expected to cause adverse effects in most breastfed infants. Two expert panels advocates waiting for at least 4 hours after a single intravenous dose of **midazolam** (e.g., for endoscopy) before resuming nursing. However, no waiting period or discarding of milk might be necessary before resuming breastfeeding after a single dose of **midazolam** in the mothers of infants over 2 months of age. [Lactmed]

Fentanyl : No waiting period or discarding of milk is required before resuming breastfeeding after **fentanyl** is used for short procedures (e.g., for endoscopy) [Lactmed]

Pethidine: A single dose for anesthesia or conscious sedation usually does not cause problems in older breastfed infants. [Lactmed]

Drugs used in this procedure are not licensed for use during breastfeeding and are therefore not recommended for use by the manufacturers. This does not imply that they are unsafe.

See information sheet on patient information leaflets on the website www.breastfeedingnetwork.org.uk for further explanation

Bibliography

British National Formulary

Hale T. W Medications in Mothers Milk 2016 (17th Ed) Hale publishing

Jones W Breastfeeding and Medication 2013 Routledge

Lactmed website <http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?LACT>

To speak to a Breastfeeding Supporter call the **National Breastfeeding Helpline 0300 100 0212**

Calls to 0300 numbers cost no more than calls to UK numbers starting 01 and 02 and will be part of any inclusive minutes that apply to your provider and call package

*The Breastfeeding Network is a Company Limited by Guarantee Registered in Scotland Company No. 330639
Registered office Whitelaw Wells, 9 Ainslie Place, Edinburgh, EH3 6AT
The Breastfeeding Network is a Registered Scottish Charity No SC027007*

©Wendy Jones PhD, MRPharmS and the Breastfeeding Network May 2017

To find your nearest Breastfeeding Supporter call the **Supporterline 0300 100 0210**



To speak to a Breastfeeding Supporter call the **National Breastfeeding Helpline 0300 100 0212**

Calls to 0300 numbers cost no more than calls to UK numbers starting 01 and 02 and will be part of any inclusive minutes that apply to your provider and call package

*The Breastfeeding Network is a Company Limited by Guarantee Registered in Scotland Company No. 330639
Registered office Whitelaw Wells, 9 Ainslie Place, Edinburgh, EH3 6AT
The Breastfeeding Network is a Registered Scottish Charity No SC027007*

©Wendy Jones PhD, MRPharmS and the Breastfeeding Network May 2017